



CLAIMS

Please amend the following claims:

1. (currently amended) A method for the preparation of a steroid modified chacotriose of general formula (Ia) or a steroid modified solatriose of general formula (Ib):




Formula (Ia)




Formula (Ib)

wherein R¹ represents a steroid or a derivative thereof selected from the group consisting of tomatidine and demissidine, and having a hydroxyl group in the 3-position and no further unprotected hydroxyl groups; and each R² independently represents a straight or branched C₁₋₁₄ alkyl group, a C₅₋₁₂ aryl or heteroaryl group optionally substituted by one or more halogen atoms or C₁₋₄ alkyl groups, or a hydroxyl group, which method comprises the step of:

reacting a compound of general formula (IIa) or (IIb)



Formula (IIa)



Formula (IIb)

wherein R³ represents a halogen atom, ~~an ethyl sulfide~~ or a sulfide group; and each R⁴ independently represents a benzoyl, substituted benzoyl, whereby the substituents are selected


from C₁₋₄ alkyl groups, halogen atoms and NO₂, acetyl or pivoly l protecting group;
with a compound of general formula (III):




Formula (III)

wherein R¹ is defined as above;

to yield a compound of general formula (IVa) or (IVb):




Formula (IVa)



Formula (IVb)

wherein R¹ and R⁴ are defined as above.

2. (original) The method according to claim 1, further comprising the step of:
deprotecting the compound of general formula (IVa) or (IVb), respectively, as defined in claim 1
to yield a compound of general formula (Va) or (Vb):



Formula (Va)




Formula (Vb)

wherein R¹ is as defined in claim 1.

3. (previously presented) The method according to claim 2 for preparing a steroid modified
chacotrose of general formula (Ia), further comprising the step of:

reacting the compound of general formula (Va) as defined in claim 2 with pivoyl chloride in the presence of an amine base to yield a compound of general formula (VIa):




Formula (VIa)

wherein R^1 represents a steroid or a derivative thereof having a hydroxyl group in the 3-position and no further unprotected hydroxyl groups, and R^5 represents a pivoyl protecting group.

4. (previously presented) The method according to claim 3 for preparing a steroid modified chactriose of general formula (Ia), further comprising the step of:

reacting the compound of general formula (VIa) as defined in claim 3 with a compound of general formula (VIIa):




Formula (VIIa)

wherein each R^2 independently represents a straight or branched C_{1-14} alkyl group, a C_{5-12} aryl or heteroaryl group optionally substituted by one or more halogen atoms or C_{1-4} alkyl groups, or a hydroxyl group;

R^3 represents a halogen atom, ~~an ethylsulfide~~ or a sulfide group; and

each R^4 independently represents a benzoyl, substituted benzoyl, whereby the substituents are selected from C_{1-4} alkyl groups, halogen atoms and NO_2 , acetyl or pivoyl protecting group;

to yield a compound general formula (VIIIa):



Formula (VIIIa)


wherein R¹, R² and R⁴ are as defined above, and R⁵ is as defined in claim 3.

5. (previously presented) The method according of claim 4 for preparing a steroid modified chacotriose of general formula (Ia), further comprising the step of:

deprotecting the compound of general formula (VIIIa) as defined in claim 4 to yield the compound of general formula (Ia).

6. (previously presented) The method according to claim 2 for preparing a steroid modified solatriose of general formula (Ib), further comprising the step of:


selectively protecting the OH groups in and 4- and 6-position of the compound of formula (Vb) as defined in claim 2 with a ketal or acetal protecting type protecting group using standard conditions, to yield a compound of general formula (VIb):



Formula (VI b)


wherein R¹ represents a steroid or a derivative thereof having a hydroxyl group in the 3-position and no further unprotected hydroxyl groups; and R⁶ represents a ketal or acetal type protecting group selected from the group consisting of benzylidene, 4-nitrobenzylidene, 4-methoxybenzylidene and isopropylidene.

7. (currently amended) The method according to claim 6 preparing a steroid modified solatriose of general formula (Ib), further comprising the step of:
reacting a compound of formula (VIb) as defined in claim 6 with a compound of general formula (VIIb):



Formula (VIIb)

wherein R³ represents a halogen atom, ~~an ethylsulfide~~ or a sulfide group; and each R⁴ independently represents a benzoyl, substituted benzoyl, whereby the substituents are selected from C₁₋₄ alkyl groups, halogen atoms and NO₂, acetyl or pivoyl protecting ~~group~~, group; to yield a compound of the general formula (VIIIb):




Formula (VIIIb)

wherein R¹ represents a steroid or a derivative thereof having a hydroxyl group in the 3-

position and no further unprotected hydroxyl groups; R⁴ is as defined above; and R⁶ is as defined in Claim 6.

8. (currently amended) The method according to claim 7 for preparing a steroid modified solatriose of general formula (Ib), further comprising the step of:

reacting a compound of formula (VIIIb) as defined in claim 7 with a compound of formula (VIIa)




Formula (VIIa)

wherein each R^2 independently represents a straight or branched C_{1-14} alkyl group, a C_{5-12} aryl or heteroaryl group optionally substituted by one or more halogen atoms or C_{1-4} alkyl groups, or a hydroxyl group;

R^3 represents a halogen atom, ~~an ethylsulfide~~ or a sulfide group; and

each R^4 independently represents a benzoyl, substituted benzoyl, whereby the substituents are selected from C_{1-4} alkyl groups, halogen atoms and NO_2 , acetyl or pivoyl protecting group;

to yield a compound of formula (IXb):



Formula (IXb)

wherein R^1 represents a steroid or a derivative thereof having a hydroxyl group in the 3-position and no further unprotected hydroxyl groups; R^2 and R^4 are as defined-above; and R^6 represents a ketal or acetal type protecting group selected from the group consisting of benzylidene, 4-nitrobenzylidene, 4-methoxybenzylidene and isopropylidene.

9. (previously presented) The method according to claim 8 for preparing a steroid modified solatriose of general formula (Ib), further comprising the step of:

deprotecting the compound of formula (IXb) as defined in claim 8 to yield the compound of formula (Ib).

10. (previously presented) The method according to claim 1, wherein R^1 represents a tomatidin-3-yl, demissidin-3-yl, solanidin-3-yl and solasodin-3-yl group.

11. (currently amended) The method according to a claim 1, wherein R^2 represents a methyl group.

12. (previously presented) The method according to claim 1, wherein R^3 in the compounds of formulae (IIa) and/or (IIb) represents a bromine atom.

13. (previously presented) The method according to claim 1, wherein the reaction is carried

out in the presence of a promoter.

14. (original) The method according to claim 13, wherein the promoter is selected from the group consisting of silver triflate, boron trifluoride diethyl etherate, trimethylsilyl triflate bromide, N-iodosuccinimide and dimethyl thiomethyl sulfonium triflate.

15. (original) The method according to claim 14, wherein the promoter is silver triflate.

16. (previously presented) The method according to claim 1, wherein the reaction is carried out under anhydrous conditions in the presence of 4Å mol sieves.

17. (previously presented) The method according to claim 2, wherein deprotection is carried out in dichloromethane or tetrahydrofuran in the presence of a C₁₋₄ alcohol and an alkali metal alkoxide having 1 to 4 carbon atoms.

18. (original) The method according to claim 17, wherein deprotection is carried out in dichloromethane in the presence of methanol and sodium methoxide.

19. (previously presented) The method according to claim 2, wherein deprotection is carried out in dichloromethane or tetrahydrofuran in the presence of water, an alkali metal hydroxide and a C₁₋₄ alcohol.

20. (original) The method according to claim 19, wherein deprotection is carried out in tetrahydrofuran, and wherein the alkali metal hydroxide is sodium hydroxide and the alcohol is methanol.

21. (original) The method according to claim 1 for preparing a steroid modified solatriose of general formula (Ib), wherein R⁴ represents a benzoyl or p-toluoyl protecting group.

22. (previously presented) The method according to claim 1, wherein reacting a compound of general formula (IIa) or (IIb) with a compound of general formula (III) is carried out in the presence of sterically hindered non-nucleophilic base.

23. (original) The method according to claim 22, wherein the sterically hindered non-nucleophilic base is selected from 2,6-lutidine, 2,4,6-collidine or 2,6-di-tertbutyl-4-methyl pyridine.

24. (previously presented) A steroid modified chacotriose of general formula (Ia) as defined in claim 1, wherein R¹ represents a tomatidin-3-yl or demissidin-3-yl group.

25. (previously presented) A compound of general formula (VIIIa) as defined in claim 4.

26. (previously presented) A compound of general formula (VIIIb) as defined in claim 7.

27. (previously presented) A compound of general formula (VIa) as defined in claim 3.


28. (previously presented) A compound of general formula (VIb) as defined in claim 6.

29. (previously presented) A compound of general formula (Va) or (Vb) as defined in claim 2.

30. (previously presented) A compound of general formula (IVa) or (IVb) as defined in claim 1.

31. (previously presented) A compound of general formula (IXb) as defined in claim 8.


32. (new) The method according to claim 1, wherein the sulfide group of R^3 is ethylsulfide.
33. (new) The method according to claim 4, wherein the sulfide group of R^3 is ethylsulfide.
34. (new) The method according to claim 7, wherein the sulfide group of R^3 is ethylsulfide.
35. (new) The method according to claim 8, wherein the sulfide group of R^3 is ethylsulfide.
36. (new) A method for the preparation a steroid modified solatriose of general formula (Ib):



Formula (Ib)

wherein R^1 represents a steroid or a derivative thereof having a hydroxyl group in the 3-position and no further unprotected hydroxyl groups; and each R^2 independently represents a straight or branched C_{1-14} alkyl group, a C_{5-12} aryl or heteroaryl group optionally substituted by one or more halogen atoms or C_{1-4} alkyl groups, or a hydroxyl group, which method comprises the step of:

reacting a compound of general formula (IIb)



Formula (IIb)

wherein R^3 represents a halogen atom or a sulfide group; and each R^4 independently represents a benzoyl, substituted benzoyl, whereby the substituents are selected from C_{1-4} alkyl groups, halogen atoms and NO_2 , acetyl or pivoyl protecting group;


with a compound of general formula (III):



Formula (III)

wherein R^1 is defined as above;

to yield a compound of general formula (IVa) or (IVb):



Formula (IVb)

wherein R^1 and R^4 are defined as above.